

Robustness of Dynamic Systems with Parameter Uncertainties

Edited by
M. Mansour
S. Balemi
W. Truöl

1992

Birkhäuser Verlag
Basel · Boston · Berlin

Contents

| | |
|--|-----|
| Preface | IX |
| Robust Stability: The Polynomial Case | |
| Kharitonov's Theorem and the Second Method of Lyapunov <i>M. Mansour and B.D.O. Anderson</i> | 3 |
| Stability Margins of Perturbed Systems <i>Y.C. Soh and Y.K. Foo</i> | 13 |
| Frequency Domain Criteria for Robust Root Locations of Generalized Disc Polynomials <i>C.B. Soh</i> | 23 |
| Computation of Value Sets of Uncertain Transfer Functions <i>W. Truöl and F.J. Kraus</i> | 33 |
| Robust Stability of General Interconnections of Interval Plants and Unstructured Norm Bounded Perturbations <i>M. Dahleh, A. Tesi and A. Vicino</i> | 43 |
| Algebraic Approach to Checking Strict Positive Real Property of Uncertain Real Rational Functions <i>T. Mori and H. Kokame</i> | 53 |
| Stability of Convex Hull of Quasipolynomials <i>V.L. Kharitonov and A.P. Zhabko</i> | 63 |
| Robust Stability: Multilinear Parameter Dependence | |
| Robustness Properties of Multilinear Interval Systems <i>H. Chapellat, L.H. Keel and S.P. Bhattacharyya</i> | 73 |
| Easily Testable Sufficient Conditions for the Robust Stability of Systems with Multiaffine Parameter Dependence <i>B.D.O. Anderson, F.J. Kraus, M. Mansour and S. Dasgupta</i> | 81 |
| Robustness Analysis for Multilinear Perturbations <i>B.T. Polyak</i> | 93 |
| On the Multiaffine Image of a Cube <i>N.-K. Tsing and A.L. Tits</i> | 105 |

| | |
|---|-----|
| Robust Stability: The Nonlinear Case | |
| Robust Absolute Stability of Continuous Systems <i>Y.Z. Tsyppkin and B.T. Polyak</i> | 113 |
| Robust Stability: The Matrix Case | |
| A Branch and Bound Method to Check the Stability of a Polytope of Matrices <i>H. Kokame and T. Mori</i> | 125 |
| Bounds on the Real Stability Radius <i>L. Qiu and E.J. Davison</i> | 139 |
| Necessary and Sufficient Conditions for Robust Stability with Structured Time-Invariant Perturbations <i>B. Bamieh and M. Dahleh</i> | 147 |
| A Nonconservative Kronecker Based Theory for Robust Root Clustering of Linear State Space Models with Real Norm Bounded Uncertainty <i>R.K. Yedavalli</i> | 155 |
| A Necessary and Sufficient Condition for Quadratic Finite Time Feedback Controllability <i>I.R. Petersen, M. Corless and E.P. Ryan</i> | 165 |
| Lyapunov Based Robustness | |
| Lyapunov Functions for Uncertain Systems <i>S. Dasgupta and G. Chockalingam</i> | 177 |
| Robustness Analysis of Linear Control Systems with Uncertain Parameters by the Method of Convex Decomposition <i>H. Kiendl and A. Michalske</i> | 189 |
| Performance | |
| H_∞ -Performance of Interval Plants and Interval Feedback Systems <i>C.V. Hollot, R. Tempo and V. Blondel</i> | 201 |
| Linear vs. Nonlinear Approaches to Robustness Analysis of LTI Control Systems <i>M. Milanese, M. Taragna, A. Trisoglio and S. Malan</i> | 211 |
| Control Design | |
| Analysis and Design of Robustly Stable Systems via Zero-Set Methods <i>E. Zehab</i> | 219 |
| A Robust Control Design in the Parameter Space <i>D.D. Šiljak</i> | 229 |
| Optimizing the Guaranteed Cost in the Control of Uncertain Linear Systems <i>I.R. Petersen and D.C. McFarlane</i> | 241 |

| | |
|---|-----|
| Robust PID Control and Lead-Lag Compensation for Linear Interval Systems | |
| <i>S.S. Ahmad, L.H. Keel and S.P. Bhattacharyya</i> | 251 |
| Model Following Robust Control of Linear Time-Varying Uncertain Systems | |
| <i>M. Fu and H. Li</i> | 261 |
| A Convex Parameterization of Robustly Stabilizing Controllers | |
| <i>A. Rantzer and A. Megretsky</i> | 271 |
| Robust Compensator Synthesis — An Experimental Application | |
| <i>G. Peretti, F. Kraus and L. Guzzella</i> | 281 |
| Velocity-Independent Yaw Eigenvalues of Four-Wheel Steering Automobiles | |
| <i>J. Ackermann</i> | 291 |
| Some Open Problems | 305 |

KHARITONOV'S THEOREM AND THE SECOND METHOD OF LYAPUNOV

Mohamed Mansour* Brian D. O. Anderson †

Abstract. In this paper Kharitonov's theorem for the robust stability of interval polynomials is proved using the second method of Lyapunov. The Hermite matrix is taken as the matrix of the quadratic form which is used as a Lyapunov function to prove Hurwitz stability. It is shown that if the four Hermite matrices corresponding to the four Kharitonov extreme polynomials are positive definite, the Hermite matrix of any polynomial of the polynomial family remains positive definite.

1. INTRODUCTION

Consider an interval polynomial

$$f(s) = \sum_{i=0}^n a_i s^i \quad (1)$$

where

$$\underline{a}_i \leq a_i \leq \bar{a}_i \quad (2)$$

Kharitonov [1] has shown that the polynomial family defined by (1) and (2) is Hurwitz stable if and only if four extreme polynomials, namely,

$$f_1(s) = \bar{a}_0 + \underline{a}_1 s + \underline{a}_2 s^2 + \bar{a}_3 s^3 + \bar{a}_4 s^4 + \underline{a}_5 s^5 + \underline{a}_6 s^6 + \dots \quad (3)$$

$$f_2(s) = \bar{a}_0 + \bar{a}_1 s + \underline{a}_2 s^2 + \underline{a}_3 s^3 + \bar{a}_4 s^4 + \bar{a}_5 s^5 + \underline{a}_6 s^6 + \dots \quad (4)$$

$$f_3(s) = \underline{a}_0 + \bar{a}_1 s + \bar{a}_2 s^2 + \underline{a}_3 s^3 + \underline{a}_4 s^4 + \bar{a}_5 s^5 + \bar{a}_6 s^6 + \dots \quad (5)$$

$$f_4(s) = \underline{a}_0 + \underline{a}_1 s + \bar{a}_2 s^2 + \bar{a}_3 s^3 + \underline{a}_4 s^4 + \underline{a}_5 s^5 + \bar{a}_6 s^6 + \dots \quad (6)$$

are Hurwitz stable.

In [2] it was shown that for 3rd degree, 4th degree and 5th degree polynomials $f(s)$, the Hurwitz stability of $f_1(s)$; $f_1(s)$ and $f_2(s)$; $f_1(s)$ and $f_2(s)$ and $f_3(s)$ respectively is necessary and sufficient for the Hurwitz stability of the interval polynomial.

Kharitonov's theorem has different proofs with the simplest one perhaps being given in [3], using the value set concept and the Cremer-Leonard-Michailov criterion.

The Hurwitz stability criterion was proved in [4], using the second method of Lyapunov with the Hermite matrix as the matrix of the quadratic form Lyapunov function. In [5] a reduced Hermite criterion is derived which makes use of a symmetric matrix of

*Automatic Control Laboratory, Swiss Federal Institute of Technology (ETH), 8092 Zurich, Switzerland, mansour@aut.ethz.ch

†Department of System Engineering, Australian National University (ANU), Canberra ACT 2600, Australia, bda101@barton.anu.edu.au

half the dimension of the original Hermite matrix and which we shall use in the sequel. The idea of the proof of Kharitonov's theorem using Lyapunov's second method is based on the following: If we assume the positive definiteness of the reduced Hermite matrix of the four Kharitonov polynomials we have to prove that the reduced Hermite matrix remains positive definite for any value of the parameters in the Kharitonov box. It is clearly sufficient to establish simply that its determinant does not go to zero. We divide the problem in two parts: We prove the above for fixed even coefficients and then for fixed odd coefficients.

The determinant of the reduced Hermite matrix is determined as a function of the even or the odd parts of the polynomial, following the line of thought in [6]. The change of the even part or the odd part of the polynomial inside the box is determined following a similar argument to that used in the proof of Markov-Chebychev theorem in [7]. In the following section we discuss two lemmas which are used in section 3 to prove Kharitonov's theorem.

2. MATHEMATICAL BACKGROUND

The polynomial $f(s)$ in (1) is partitioned into its even and odd parts as follows:

$$f(s) = \sum_{i=0}^n a_i s^i = h(s^2) + sg(s^2) \quad (7)$$

The four Kharitonov polynomials are given (in obvious notation) by:

$$\begin{aligned} f_1 &= (\bar{h}, g) \\ f_2 &= (\bar{h}, \bar{g}) \\ f_3 &= (\underline{h}, \bar{g}) \\ f_4 &= (\underline{h}, g) \end{aligned} \quad (8)$$

The $n \times n$ Hermite matrix P associated with $f(s)$ is given by:

$$p_{ij} = \begin{cases} \sum_{k=1}^i (-1)^{k+1} a_{n-k+1} a_{n-i-j+k} & j \geq i, j+i \text{ even} \\ p_{ji} & j < i, j+i \text{ even} \\ 0 & j+i \text{ odd} \end{cases} \quad (9)$$

For $n = 6$ for example,

$$P = \begin{bmatrix} a_6 a_5 & 0 & a_6 a_3 & 0 & a_6 a_1 & 0 \\ 0 & -a_6 a_3 + a_5 a_4 & 0 & -a_6 a_1 + a_5 a_2 & 0 & a_5 a_0 \\ a_6 a_3 & 0 & a_6 a_1 - a_5 a_2 + a_4 a_3 & 0 & -a_5 a_0 + a_4 a_1 & 0 \\ 0 & -a_6 a_1 + a_5 a_2 & 0 & a_5 a_0 - a_4 a_1 + a_3 a_2 & 0 & a_3 a_0 \\ a_6 a_1 & 0 & -a_5 a_0 + a_4 a_1 & 0 & -a_3 a_0 + a_2 a_1 & 0 \\ 0 & a_5 a_0 & 0 & a_3 a_0 & 0 & a_1 a_0 \end{bmatrix} \quad (10)$$

For simplicity and without real loss of generality we shall assume that $a_n = 1$. The Hermite criterion states that $f(s)$ is Hurwitz stable if and only if P is positive definite. Consider the matrices

$$A = \begin{bmatrix} 0 & 1 & & \\ \vdots & \vdots & \ddots & \\ 0 & 0 & \cdots & 1 \\ -a_0 & -a_1 & \cdots & -a_{n-1} \end{bmatrix}, \quad J = \begin{bmatrix} 0 & \cdots & 0 & 1 \\ \vdots & & & 0 \\ 0 & & & \vdots \\ 1 & 0 & \cdots & 0 \end{bmatrix}, \quad P^* = JPJ \quad (11)$$

Then we get the Lyapunov equation

$$A^T P^* + P^* A = -2\underline{b} \underline{b}^T \quad (12)$$

where

$$\underline{b}^T = [\dots 0 \ a_{n-3} \ 0 \ a_{n-1}]$$

and $\underline{x}^T P^* \underline{x}$ is the Lyapunov function which proves the Hurwitz stability of $f(s)$.

The reduced Hermite matrices C and D derived in [5] are submatrices of the Hermite matrix P obtainable by deleting even-numbered rows and columns from P or deleting odd-numbered rows and columns respectively.

For $n = 6$ for example

$$C = \begin{bmatrix} a_6 a_5 & & a_6 a_3 & & a_6 a_1 \\ a_6 a_3 & a_6 a_1 - a_5 a_2 + a_4 a_3 & & -a_5 a_0 + a_4 a_1 \\ a_6 a_1 & -a_5 a_0 + a_4 a_1 & & -a_3 a_0 + a_2 a_1 \end{bmatrix} \quad (13)$$

$$D = \begin{bmatrix} -a_6 a_3 + a_5 a_4 & -a_6 a_1 + a_5 a_2 & a_5 a_0 \\ -a_6 a_1 + a_5 a_2 & a_5 a_0 - a_4 a_1 + a_3 a_2 & a_3 a_0 \\ a_5 a_0 & a_3 a_0 & a_1 a_0 \end{bmatrix} \quad (14)$$

A form of the result in [5] can be stated as follows:

If $a_i > 0 \quad i = 0, 1, \dots, n$ then

$$P > 0 \Leftrightarrow C > 0 \Leftrightarrow D > 0 \quad (15)$$

If (with $a_n = 1$)

$$A_1 = \begin{bmatrix} 0 & 1 & & \\ \vdots & 0 & 1 & \\ \vdots & \vdots & 0 & \ddots \\ \vdots & \vdots & \vdots & \\ 0 & 0 & 0 & 1 \\ -a_0 & -a_2 & -a_4 & -a_{n-2} \end{bmatrix} \quad (16)$$

then there holds

$$(JCJ)A_1 = A_1^T(JCJ) = -JDJ \quad (17)$$

(where J is as above except for its dimension).
By way of digression we may recall [5], [8] that

$$\begin{aligned} \text{for } n \text{ even: } \quad \det C &= \det H_{n-1}, \quad \det D = a_0 \det H_{n-1} \\ \text{for } n \text{ odd: } \quad \det C &= a_0 \det H_{n-1}, \quad \det D = \det H_{n-1} \end{aligned} \quad (18)$$

where $\det H_{n-1}$ is the $n-1$ Hurwitz-determinant. H_n is given by

$$H_n = \begin{bmatrix} a_1 & a_0 & 0 & \cdots & \cdots & \cdots & \cdots & 0 \\ a_3 & a_2 & a_1 & a_0 & \cdots & \cdots & \cdots & 0 \\ a_5 & a_4 & a_3 & a_2 & a_1 & a_0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & a_{n-2} \\ 0 & 0 & \cdots & \cdots & \cdots & \cdots & \cdots & a_n \end{bmatrix} \quad (19)$$

It is clear that if C or $D > 0$ then $\det H_{n-1}$ is positive. The quantity $\det H_{n-1}$ determines the critical stability condition. If coefficients of an originally stable system are perturbed, stability will be lost if $\det C$ (or $\det D$) becomes zero.

Without loss of generality we shall consider in the sequel even polynomials by setting $n = 2m$. A similar treatment can be used for odd n .

Lemma 1 *With notation as above,*

$$\det C = (-1)^{\frac{m(m-1)}{2}} \prod g(\alpha_i) = (-1)^{\frac{m(m-1)}{2}} a_{n-1} \prod h(\beta_j) \quad (20)$$

where α_i and β_j are the roots of $h(\lambda)$ and $g(\lambda)$ respectively.

Proof: The proof follows the same lines as [6]. Consider the transfer function

$$w_1(\lambda) = \frac{g(\lambda)}{h(\lambda)} = \frac{a_1 + a_3\lambda + \dots + a_{n-1}\lambda^{m-1}}{a_0 + a_2\lambda + \dots + \lambda^m} \quad (21)$$

A realisation of (21) is given by

$$A_1 = \begin{bmatrix} 0 & 1 & & & \\ 0 & & \ddots & & \\ & & & 1 & \\ -a_0 & -a_2 & \cdots & -a_{n-2} & \end{bmatrix}, \quad b_1 = \begin{bmatrix} 0 \\ \vdots \\ 0 \\ 1 \end{bmatrix}, \quad c_1^T = [a_1 \ a_3 \ \dots \ a_{n-1}] \quad (22)$$

Define also

$$C^* = JCJ \quad D^* = JDJ \quad (23)$$

The observability matrix is given by

$$S_1 = \begin{bmatrix} c_1^T \\ c_1^T A_1 \\ \vdots \\ c_1^T A_1^{m-1} \end{bmatrix} = S_1(A_1, c_1^T) \quad (24)$$

Let e_1, e_2, \dots, e_m be the unit vectors; then it is easily verified that

$$S_1(A_1, e_1^T) = I, S_1(A_1, e_2^T) = A_1, \dots, S_1(A_1, e_m^T) = A_1^{m-1}$$

Therefore,

$$S_1(A_1, e^T) = a_1 I + a_3 A_1 + \dots + a_{n-1} A_1^{m-1} = g(A_1) \quad (25)$$

But

$$C^* e_m = \begin{bmatrix} a_1 \\ a_3 \\ \vdots \\ a_{n-3} \\ a_{n-1} \end{bmatrix} = e_1$$

and by (17)

$$C^* A_1^k = (A_1^T)^k C^*, k = 1, 2, 3, \dots$$

Then it can be seen by inspection that

$$C^* [e_m \ A_1 e_m \ \dots \ A_1^{m-1} e_m] = S_1^T \quad (26)$$

But $[e_m \ A_1 e_m \ \dots \ A_1^{m-1} e_m]$ is clearly a lower triangular matrix with second diagonal elements equal to one.

Therefore, $\det C^* = (-1)^{\frac{m(m-1)}{2}} \det S_1^T$ and from (25)

$$\det C^* = (-1)^{\frac{m(m-1)}{2}} \det g(A_1) \quad (27)$$

From a well known result in matrix theory $\det g(A_1) = \prod g(\alpha_i)$ where α_i are the eigenvalues of A_1 .

Therefore,

$$\det C = (-1)^{\frac{m(m-1)}{2}} \prod g(\alpha_i) \quad (28)$$

where α_i are the roots of $h(\lambda)$.

For the proof of the second equality of lemma 1, consider the transfer function

$$w_2(\lambda) = \frac{h(\lambda)}{\lambda^2 g(\lambda)} = \frac{a_0 + a_2 \lambda + \dots + \lambda^m}{\frac{a_1}{a_{n-1}} \lambda^2 + \frac{a_3}{a_{n-1}} \lambda^3 + \dots + \lambda^{m+1}} \quad (29)$$

A realisation of (29) is given by

$$A_2 = \begin{bmatrix} 0 & 1 & & & \\ \vdots & 0 & 1 & & \\ \vdots & \vdots & & \ddots & 1 \\ 0 & 0 & -\frac{a_1}{a_{n-1}} & \dots & -\frac{a_{n-3}}{a_{n-1}} \end{bmatrix}, \quad b_2 = \begin{bmatrix} 0 \\ \vdots \\ 0 \\ 1 \end{bmatrix}, \quad e_2^T = [a_0 \ a_2 \ \dots \ 1] \quad (30)$$

B

The observability matrix S_2 is given by

$$S_2 = \begin{bmatrix} \underline{e}_2^T \\ \underline{e}_2^T A_2 \\ \vdots \\ \underline{e}_2^T A_2^m \end{bmatrix} = \begin{bmatrix} a_0 & a_2 & \cdots & 1 \\ 0 & a_0 & \cdots & \cdots \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & a_0 + \dots \end{bmatrix} = S_2(A_2, \underline{e}_2^T) \quad (31)$$

$h(B)$ (established below)

Now

$$S_2(A_2, \underline{e}_1^T) = I, \quad S_2(A_2, \underline{e}_2^T) = A_2, \dots, S_2(A_2, \underline{e}_{m+1}^T) = A_2^m, \quad \text{so that} \\ S_2(A_2, \underline{e}_2^T) = a_0 I + a_2 A_2 + \dots + A_2^m = h(A_2) \quad (32)$$

If we denote the $m \times m$ submatrix of A_2 obtained by deleting the first row and column by B then the corresponding submatrix of S_2 is evidently $h(B)$.

We have

$$\det h(A) = a_0 \det h(B) \quad (33)$$

Now it can be easily established by direct calculation that

$$D^* B = B^T D^* \quad (34)$$

and

$$D^* \underline{e}_m = a_{n-1} \begin{bmatrix} a_0 \\ a_2 \\ \vdots \\ \vdots \\ a_{n-2} \end{bmatrix} - \begin{bmatrix} 0 \\ a_1 \\ a_3 \\ \vdots \\ a_{n-3} \end{bmatrix} = \underline{f} \quad (35)$$

Also

$$\underline{e}_2^T A_2 = \frac{1}{a_{n-1}} \begin{bmatrix} 0 & \underline{f}^T \end{bmatrix} \quad (36)$$

Recalling the construction of S_2 above and its submatrix $h(B)$, we see that

$$\begin{bmatrix} \underline{f} & B \underline{f} \dots B^{m-1} \underline{f} \end{bmatrix} = a_{n-1} [h(B)]^T \quad (37)$$

accordingly

$$D^* \begin{bmatrix} \underline{e}_m \dots B^{m-1} \underline{e}_m \end{bmatrix} = \begin{bmatrix} D^* \underline{e}_m & B^T D^* \underline{e}_m \dots (B^T)^{m-1} D^* \underline{e}_m \end{bmatrix} = a_{n-1} [h(B)]^T \quad (38)$$

Now $\begin{bmatrix} \underline{e}_m \dots B^{m-1} \underline{e}_m \end{bmatrix}$ is a lower triangular matrix with unity elements in the second diagonal, therefore,

$$\det D^* = (-1)^{\frac{m(m-1)}{2}} a_{n-1} \det h(B) = (-1)^{\frac{m(m-1)}{2}} a_{n-1} h(0) \prod h(\beta_j) \\ = (-1)^{\frac{m(m-1)}{2}} a_0 a_{n-1} \prod h(\beta_j) \quad (39)$$

where β_j are the roots of $g(\lambda)$. Hence

$$\det C = \frac{1}{a_0} \det D^* = (-1)^{\frac{m(m-1)}{2}} a_{n-1} \prod h(\beta_j)$$

The second key result which we need is as follows:

Lemma 2 *With quantities as defined above*

$$\lambda^T C^* \lambda = g(\lambda) \frac{dh(\lambda)}{d\lambda} - h(\lambda) \frac{dg(\lambda)}{d\lambda} \quad (40)$$

where $\lambda^T = [1 \ \lambda \ \dots \ \lambda^{m-1}]$

Proof: It is easily verified that

$$\frac{h(\mu)g(\lambda) - h(\lambda)g(\mu)}{\mu - \lambda} = [1 \ \mu \dots \mu^{m-1}] C^* \begin{bmatrix} 1 \\ \lambda \\ \vdots \\ \lambda^{m-1} \end{bmatrix} \quad (41)$$

It is straightforward to show that if we substitute $\mu = \lambda + \varepsilon$ and let ε tends to zero

$$\frac{h(\lambda + \varepsilon)g(\lambda) - h(\lambda)g(\lambda + \varepsilon)}{\varepsilon} = \frac{[h(\lambda + \varepsilon) - h(\lambda)]g(\lambda) - [g(\lambda + \varepsilon) - g(\lambda)]h(\lambda)}{\varepsilon}$$

$$\implies g(\lambda) \frac{dh(\lambda)}{d\lambda} - h(\lambda) \frac{dg(\lambda)}{d\lambda}$$

Therefore,

$$\lambda^T C^* \lambda = g(\lambda) \frac{dh(\lambda)}{d\lambda} - h(\lambda) \frac{dg(\lambda)}{d\lambda} \quad (42)$$

3. PROOF OF KHARITONOV'S THEOREM

We need the following additional lemma:

Lemma 3 *Consider the family of polynomials defined by*

$$f(s) = \sum_{i=0}^n a_i s^i$$

$$\text{and } \underline{a}_{2j+1} \leq a_{2j+1} \leq \bar{a}_{2j+1} \quad \text{for } j = 0, 1, \dots, m-1$$

while a_{2j} is fixed at the values of \bar{h} where $\bar{h} = \{\underline{h}, \bar{h}\}$. Suppose that C is positive definite for the four Kharitonov polynomials. Then $\det C$ is nonzero for all polynomials in the family.

Proof: Let $a_i = (-1)^{\frac{i-1}{2}} \varphi_i(t)$ for $i = 1, 3, 5, \dots, n-1$ where $\varphi_i(t)$ are monotonically increasing functions and differentiable when t varies from $t = t_1$ to $t = t_2$. Taking $t = t_1$ gives g and taking $t = t_2$ gives \bar{g} . Then

$$\begin{aligned} \frac{dg}{dt} &= \sum (-1)^{\frac{i-1}{2}} \frac{\partial g}{\partial a_i} \frac{d\varphi_i}{dt} \\ &= \frac{d\varphi_1}{dt} - \lambda \frac{d\varphi_3}{dt} + \lambda^2 \frac{d\varphi_5}{dt} - \dots \end{aligned} \quad (43)$$

At the roots α_i of $\bar{h}(\lambda)$, which are negative and independent of t , we get

$$\frac{dg(\alpha_i)}{dt} > 0 \quad (44)$$

Now from lemma 2 and the stability of the four Kharitonov polynomials

$$\bar{g}(\alpha_i) \frac{d\bar{h}(\alpha_i)}{d\alpha_i} > 0 \quad \text{and} \quad \underline{g}(\alpha_i) \frac{d\bar{h}(\alpha_i)}{d\alpha_i} > 0 \quad (45)$$

which means that $g(\alpha_i)$ does not change sign as t varies from t_1 to t_2 . Then from lemma 1 $\det C$ does not go to zero. Hence C remains positive definite for all values of a_1, a_3, \dots, a_{n-1} in the given range. ■

Now let us return to the family defined by (1) and (2), with the assumption that the four Kharitonov polynomials are Hurwitz stable. By lemma 3, if we choose any $g(\lambda)$ consistent with (2), the matrix C associated with $g(\lambda)$ and either $\bar{h}(\lambda)$ or $\underline{h}(\lambda)$ is positive definite. Now let $a_i = (-1)^{\frac{i}{2}} \varphi_i(t)$ $i = 0, 2, 4, \dots, n-2$ where $\varphi_i(t)$ are monotonically increasing and differentiable when t varies from $t = t_1$ to $t = t_2$. Taking $t = t_1$ gives \underline{h} and taking $t = t_2$ gives \bar{h} . Then

$$\begin{aligned} \frac{d\bar{h}}{dt} &= \sum (-1)^{\frac{i}{2}} \frac{\partial \bar{h}}{\partial a_i} \frac{d\varphi_i}{dt} \\ &= \frac{d\varphi_0}{dt} - \lambda \frac{d\varphi_2}{dt} + \lambda^2 \frac{d\varphi_4}{dt} - \dots \end{aligned} \quad (46)$$

and at all roots β_j of $g(\lambda)$ which are negative we get

$$\frac{dh(\beta_j)}{dt} > 0 \quad (47)$$

Then from lemma 2

$$-\bar{h}(\beta_j) \frac{dg(\beta_j)}{d\beta_j} > 0 \quad \text{and} \quad -\underline{h}(\beta_j) \frac{dg(\beta_j)}{d\beta_j} > 0 \quad (48)$$

which means that $h(\beta_j)$ does not change sign as t varies from t_1 to t_2 . Then from lemma 1 $\det C$ does not go to zero. Hence C remains positive definite for all values of a_i in the given range for $i = 0, 1, \dots, n$. Therefore, if C is positive definite for the four Kharitonov polynomials it remains positive definite for the whole box.

4. CONCLUSIONS

It was shown that the positive definiteness of the Hermite matrix for the four Kharitonov polynomials guarantees its positive definiteness for all other values of the coefficients inside the Kharitonov box.

As the Hermite matrix can be used to construct a Lyapunov function to prove Hurwitz stability, then the above result can be considered as a Lyapunov-Kharitonov link.

Nowhere in our argument have we used the concept of a value set, nor monotony of the argument of a Hurwitz stable polynomial evaluated on the imaginary axis. This monotony can actually be easily derived by using a property like that of lemma 2, but applied to the full size Hermite matrix P , rather than C .

REFERENCES

- [1] V. L. Kharitonov. Asymptotic stability of an equilibrium position of a family of systems of linear differential equations. *Differentsial'nye Uravneniya*, 14:2086-2088, 1978.
- [2] B. D. O. Anderson, E. I. Jury, and M. Mansour. On robust Hurwitz polynomials. *IEEE Transactions on Automatic Control*, 32:909-913, 1987.
- [3] R. J. Minnichelli, J. J. Anagnost, and C. A. Desoer. An elementary proof of Kharitonov's stability theorem with extensions. *IEEE Transactions on Automatic Control*, 34(9):995-998, 1989.
- [4] P. C. Parks. A new proof of the Routh-Hurwitz stability criterion using the second method of Lyapunov. *Proceedings of the Cambridge Phil. Soc.*, 58:694-702, 1962.
- [5] B. D. O. Anderson. The reduced Hermite criterion with application to proof of the Liénard-Chipard criterion. *IEEE Transactions on Automatic Control*, 17:669-672, 1972.
- [6] R. E. Kalman. Mathematical description of linear dynamical systems. *J. SIAM Control, Ser. A*, 1(2):152-192, 1963.
- [7] F. R. Gantmacher. *The theory of matrices*. Chelsea Publ. Co., New York, 2 edition, 1960.
- [8] E. I. Jury. Inners and stability of dynamic systems. *R. E. Krieger Publ. Co., Florida*, 1982.